

Guidelines for Perioperative Care for Pancreaticoduodenectomy: Enhanced Recovery After Surgery (ERAS[®]) Society Recommendations

Kristoffer Lassen · Marielle M. E. Coolson · Karem Slim · Francesco Carli · José E. de Aguilar-Nascimento · Markus Schäfer · Rowan W. Parks · Kenneth C. H. Fearon · Dileep N. Lobo · Nicolas Demartines · Marco Braga · Olle Ljungqvist · Cornelis H. C. Dejong

© Société Internationale de Chirurgie 2012

Abstract

Background Protocols for enhanced recovery provide comprehensive and evidence-based guidelines for best perioperative care. Protocol implementation may reduce complication rates and enhance functional recovery and, as a result of this, also reduce length-of-stay in hospital. There is no comprehensive framework available for pancreaticoduodenectomy.

Methods An international working group constructed within the Enhanced Recovery After Surgery (ERAS[®]) Society constructed a comprehensive and evidence-based framework for best perioperative care for pancreaticoduodenectomy

This work was conducted on behalf of ERAS[®] Society, the European Society for Clinical Nutrition and Metabolism and the International Association for Surgical Metabolism and Nutrition.

The guidelines are published as a joint effort between the Enhanced Recovery After Surgery (ERAS) Society, for Perioperative Care, The European Society for Clinical Nutrition and Metabolism (ESPEN) and The International Association for Surgical Metabolism and Nutrition (IASMEN) and copyrights for this publication is shared between the three societies. The guidelines are published jointly in World Journal of Surgery (IASMEN) and Clinical Nutrition (ESPEN), and will also be available on the ESPEN (<http://www.espen.org>) and ERAS Society website (<http://www.erassociety.org>).

K. Lassen (✉)

Department of GI and HPB Surgery, University Hospital Northern Norway, Tromsø and Institute of Clinical Medicine, University of Tromsø, Tromsø, Norway
e-mail: lassen@unn.no; xtofero@gmail.com

K. Lassen · R. W. Parks · K. C. H. Fearon
Clinical Surgery, University of Edinburgh, Royal Infirmary of Edinburgh, Edinburgh, United Kingdom

M. M. E. Coolson · C. H. C. Dejong
Department of Surgery, University Hospital Maastricht and NUTRIM School for Nutrition, Toxicology and Metabolism, Maastricht, The Netherlands

patients. Data were retrieved from standard databases and personal archives. Evidence and recommendations were classified according to the GRADE system and reached through consensus in the group. The quality of evidence was rated “high”, “moderate”, “low” or “very low”. Recommendations were graded as “strong” or “weak”.

Results Comprehensive guidelines are presented. Available evidence is summarised and recommendations given for 27 care items. The quality of evidence varies substantially and further research is needed for many issues to improve the strength of evidence and grade of recommendations.

Conclusions The present evidence-based guidelines provide the necessary platform upon which to base a unified protocol for perioperative care for pancreaticoduodenectomy. A unified protocol allows for comparison between centres and across national borders. It facilitates multi-institutional prospective cohort registries and adequately powered randomised trials.

Introduction

Enhanced Recovery After Surgery (ERAS), Fast-Track or Clinical Pathway programmes are multimodal strategies that

K. Slim

Department of Digestive Surgery, Hospital CHU Estaing, Clermont-Ferrand, France

F. Carli

Department of Anesthesia, McGill University Health Centre, Montreal, QC, Canada

J. E. de Aguilar-Nascimento

Department of Surgery, Federal University of Mato Grosso, Cuiaba, Brazil

aim to attenuate the loss of, and improve the restoration of, functional capacity after surgery. Morbidity is reduced [1] and recovery enhanced by reducing surgical stress, by optimal control of pain, early oral diet and early mobilisation. As a consequence, length-of-stay in hospital (LOSH) and costs are also reduced. The ERAS group has published evidence-based consensus recommendations for colorectal surgery [2, 3]. Beneficial experiences with clinical pathway programmes after pancreaticoduodenectomy (PD, Whipple's procedure) have been published [4–9], but the reported series employed different protocols, or no prospective protocol at all [6]. A comprehensive consensus framework is presented on which to base a future protocol for optimal perioperative care after PD. Such a recommendation will allow for a unified protocol to be developed and validated prospectively across different institutions and healthcare systems. This guideline framework has been formulated and endorsed by the ERAS Society, European Society for Clinical Nutrition and Metabolism (ESPEN) and the International Association for Surgical Metabolism and Nutrition (IASMEN).

Methods

Literature search

The authors met in April 2011 and the topics to be included were agreed and allocated. A principal literature search up to June 2011 was undertaken. Comprehensive drafts were circulated for discussion and reviewed in a group conference in November 2011. Additional relevant literature published after June 2011 was considered by members of the group at meetings in November 2011 and May 2012.

Study selection

All co-authors screened web-based databases and personal archives for relevant articles. Non-systematic emphasis was given to more recent publications and publications of better quality (moderate- and high-quality randomised controlled trials and high-quality, large cohort studies; and systematic reviews and meta-analyses of these). Retrospective series were considered only if data of better quality could not be identified.

M. Schäfer · N. Demartines
Department of Visceral Surgery, University Hospital of
Lausanne (CHUV), Lausanne, Switzerland

D. N. Lobo
Division of Gastrointestinal Surgery, Nottingham Digestive
Diseases Centre National Institute for Health Research
Biomedical Research Unit, Nottingham University Hospitals,
Queen's Medical Centre, Nottingham, United Kingdom

M. Braga
San Raffaele University, Milan, Italy

Quality assessment and grading

The strength of evidence and conclusive recommendations were assessed and agreed by all authors in May 2012. Quality of evidence and recommendations were evaluated according to the Grading of Recommendations, Assessment, Development and Evaluation (GRADE) system [10–12]. Quoting from the GRADE guidelines [12], the recommendations are: “Strong recommendations indicate that the panel is confident that the desirable effects of adherence to a recommendation outweigh the undesirable effects”. “Weak recommendations indicate that the desirable effects of adherence to a recommendation probably outweigh the undesirable effects, but the panel is less confident”. Recommendations are based on quality of evidence (high, moderate, low, very low) but also on the balance between desirable and undesirable effects; and on values and preferences [12]. The latter implies that, in some cases, strong recommendations may be reached from low-quality data and vice versa. A summary of the guidelines is shown in Table 1.

Evidence and recommendations

Preoperative counselling

Preoperative counselling targeting expectations about surgical and anaesthetic procedures may diminish fear and anxiety and enhance postoperative recovery and discharge [13–15]. Personal counselling, leaflets or multimedia information containing explanations of the procedure along with tasks that the patient should be encouraged to fulfil may improve perioperative feeding, early postoperative mobilisation, pain control, and respiratory physiotherapy; and hence reduce the risk of complications [16–18]. Ideally, the patient should meet with the surgeon, anaesthetist and nurse.

Summary and recommendation	Patients should receive dedicated preoperative counselling routinely.
Evidence level	Low
Recommendation grade	Strong

O. Ljungqvist
Department of Surgery, Örebro University Hospital, Örebro,
Sweden

O. Ljungqvist
Department of Molecular Medicine and Surgery, Karolinska
Institutet, Stockholm, Sweden

Table 1 Guidelines for perioperative care for pancreaticoduodenectomy: Enhanced Recovery After Surgery (ERAS[®]) Society recommendations

Item	Summary and recommendations	Evidence level	Recommendation grade
Preoperative counselling	Patients should receive dedicated preoperative counselling routinely	Low	Strong
Perioperative biliary drainage	Preoperative endoscopic biliary drainage should not be undertaken routinely in patients with a serum bilirubin concentration <250 µmol/l	Moderate	Weak
Preoperative smoking and alcohol consumption	For alcohol abusers, 1 month of abstinence before surgery is beneficial and should be attempted. For daily smokers, 1 month of abstinence before surgery is beneficial. For appropriate groups, both should be attempted	Alcohol abstinence: low Smoking cessation: moderate	Strong
Preoperative nutrition	Routine use of preoperative artificial nutrition is not warranted, but significantly malnourished patients should be optimized with oral supplements or enteral nutrition preoperatively	Very low	Weak
Perioperative oral immunonutrition (IN)	The balance of evidence suggests that IN for 5–7 days perioperatively should be considered because it may reduce the rate of infectious complications in patients undergoing major open abdominal surgery	Moderate	Weak
Oral bowel preparation	Extrapolation of data from studies on colonic surgery and retrospective studies in PD show that MBP has no proven benefit. MBP should not be used	Moderate	Strong
Preoperative fasting and preoperative treatment with carbohydrates	Intake of clear fluids up to 2 h before anaesthesia does not increase gastric residual volume and is recommended before elective surgery. Intake of solids should be withheld 6 h before anaesthesia. Data extrapolation from studies in major surgery suggests that preoperative oral carbohydrate treatment should be given in patients without diabetes	Fluid intake: high Solid intake: low Carbohydrate loading: low	Fasting: strong carbohydrate loading: strong
Preanaesthetic medication	Data from studies on abdominal surgery show no evidence of clinical benefit from pre-operative use of long-acting sedatives, and they should not be used routinely. Short-acting anxiolytics may be used for procedures such as insertion of epidural catheters	No long-acting sedatives: moderate	Weak
Anti-thrombotic prophylaxis	LMWH reduces the risk of thromboembolic complications, and administration should be continued for 4 weeks after hospital discharge. Concomitant use of epidural analgesia necessitates close adherence to safety guidelines. Mechanical measures should probably be added for patients at high risk	High	Strong
Antimicrobial prophylaxis and skin preparation	Antimicrobial prophylaxis prevents surgical-site infections, and should be used in a single-dose manner initiated 30–60 min before skin incision. Repeated intraoperative doses may be necessary depending on the half-life of the drug and duration of procedure	High	Strong
Epidural analgesia	Mid-thoracic epidurals are recommended based on data from studies on major open abdominal surgery showing superior pain relief and fewer respiratory complications compared with intravenous opioids	Pain: high Reduced respiratory complications: moderate Overall morbidity: low	Weak
Intravenous analgesia	Some evidence supports the use of PCA or intravenous lidocaine analgesic methods. There is insufficient information on outcome after PD	PCA: very low I.V. Lidocaine: moderate	Weak
Wound catheters and transversus abdominis plane block	Some evidence supports the use of wound catheters or TAP blocks in abdominal surgery. Results are conflicting and variable, and mostly from studies on lower gastrointestinal surgery	Wound catheters: moderate TAP blocks: moderate	Weak

Table 1 continued

Item	Summary and recommendations	Evidence level	Recommendation grade
Postoperative nausea and vomiting (PONV)	Data from the literature on gastrointestinal surgery in patients at risk of PONV show the benefits of using different pharmacological agents depending on the patient's PONV history, type of surgery and type of anaesthesia. Multimodal intervention during and after surgery is indicated	Low	Strong
Incision	The choice of incision is at the surgeon's discretion, and should be of a length sufficient to ensure good exposure	Very low	Strong
Avoiding hypothermia	Intraoperative hypothermia should be avoided by using cutaneous warming, i.e., forced-air or circulating-water garment systems	High	Strong
Postoperative glycaemic control	Insulin resistance and hyperglycaemia are strongly associated with postoperative morbidity and mortality. Treatment of hyperglycaemia with intravenous insulin in the ICU setting improves outcomes but hypoglycaemia remains a risk. Several ERAS protocol items attenuate insulin resistance and facilitate glycaemic control without the risk of hypoglycaemia. Hyperglycaemia should be avoided as far as possible without introducing the risk of hypoglycaemia	Low	Strong
Nasogastric intubation	Pre-emptive use of nasogastric tubes postoperatively does not improve outcomes, and their use is not warranted routinely	Moderate	Strong
Fluid balance	Near-zero fluid balance, avoiding overload of salt and water results in improved outcomes. Perioperative monitoring of stroke volume with transoesophageal Doppler to optimize cardiac output with fluid boluses improves outcomes. Balanced crystalloids should be preferred to 0.9 % saline	Fluid balance: high oesophageal doppler: moderate Balanced crystalloids vs. 0.9 % saline: moderate	Strong
Perianastomotic drain	Early removal of drains after 72 h may be advisable in patients at low risk (i.e., amylase content in drain <5,000 U/L) for developing a pancreatic fistula. There is insufficient evidence to recommend routine use of drains, but their use is based only on low-level evidence	Early removal: high	Early removal: strong
Somatostatin analogues	Somatostatin and its analogues have no beneficial effects on outcome after PD. In general, their use is not warranted. Subgroup analyses for variability in the texture and duct size of the pancreas are not available	Moderate	Strong
Urinary drainage	Suprapubic catheterisation is superior to transurethral catheterisation if used for >4 days. Transurethral catheters can be removed safely on postoperative day 1 or 2 unless otherwise indicated	High	For suprapubic: weak Transurethral catheter out POD 1–2: strong
Delayed gastric emptying (DGE)	There are no acknowledged strategies to avoid DGE. Artificial nutrition should be considered selectively in patients with DGE of long duration	Very low	Strong
Stimulation of bowel movement	A multimodal approach with epidural and near-zero fluid balance is recommended. Oral laxatives and chewing gum given postoperatively are safe, and may accelerate gastrointestinal transit	Laxatives: very low Chewing gum: low	Weak
Postoperative artificial nutrition	Patients should be allowed a normal diet after surgery without restrictions. They should be cautioned to begin carefully and increase intake according to tolerance over 3–4 days. Enteral tube feeding should be given only on specific indications and parenteral nutrition should not be employed routinely	Early diet at will: moderate	Strong
Early and scheduled mobilisation	Patients should be mobilized actively from the morning of the first postoperative day and encouraged to meet daily targets for mobilisation	Very low	Strong
Audit	Systematic improves compliance and clinical outcomes	Low	Strong

Preoperative biliary drainage

Five meta-analyses [19–23], and two articles from a randomised controlled trial (RCT) not included in the meta-analyses [24, 25], assessed the role of biliary drainage before PD. The first meta-analysis from 2002 [19], included randomised ($n = 5$) and non-randomised trials ($n = 18$). A Cochrane review [21] included 5 randomised trials, but considered all 5 trials to have a risk of bias, thereby weakening the conclusions reached. Of the trials included, 4 evaluated percutaneous drainage and 1 evaluated endoscopic drainage. The Cochrane review concluded that preoperative biliary drainage did not decrease mortality in patients with obstructive jaundice. Although there was a trend towards decreased postoperative morbidity, the increased risk of procedure-related complications counterbalanced this possible benefit (especially for percutaneous drainage). The findings of the Cochrane review were in accordance with those of the other meta-analyses, suggesting that preoperative drainage confers neither benefit nor harm. One recent RCT not included in the meta-analyses [24] (and which included patients with serum bilirubin concentrations $<250 \mu\text{mol/l}$) showed increased morbidity in patients undergoing preoperative biliary drainage (endoscopic primarily; percutaneous as rescue option), but the delay in surgery did not affect overall survival [25].

Summary and recommendation	Preoperative endoscopic biliary drainage should not be carried out routinely in patients with a serum bilirubin concentration $<250 \mu\text{mol/l}$
Evidence level	Moderate
Recommendation grade	Weak

Preoperative smoking and alcohol consumption

Overall postoperative morbidity is increased by two- to threefold in alcohol abusers [26]. Also, 1 month of preoperative abstinence has been shown to significantly improve outcome in a group who took “five or more drinks (60 g of ethanol) a day without clinical or historical evidence of alcohol related illness” [27].

Daily smokers (>2 cigarettes daily for ≥ 1 year) have an increased risk of pulmonary and wound complications [28, 29]. RCTs have demonstrated reductions in the rates of both types of complications 1 month after cessation of smoking [29, 30].

Summary and recommendation	For alcohol abusers, 1 month of abstinence before surgery is beneficial and
----------------------------	---

should be attempted. For daily smokers, 1 month of abstinence before surgery is beneficial. For appropriate groups, both should be attempted

Evidence level	Alcohol abstinence: Low; Smoking cessation: Moderate
----------------	--

Recommendation grade	Strong
----------------------	--------

Preoperative nutrition

In western countries, patients scheduled for PD are, in general, not malnourished, and usually present with $<7\%$ weight loss [31]. In such cases, preoperative artificial nutrition is not warranted. The situation may be different in other regions. It is widely accepted that significantly malnourished patients suffer increased postoperative morbidity after major surgery [32–34]. Preoperative supplements with oral sip feeds or enteral tube feeds are usually administered in these cases, but scientific evidence to support this routine (as opposed to no nutritional support) is lacking. Extrapolating data from studies in the postoperative setting suggests that parenteral nutrition should be used only if the enteral route is inaccessible.

Summary and recommendation	Routine use of preoperative artificial nutrition is not warranted, but significantly malnourished patients should be optimized with oral supplements or enteral nutrition preoperatively
----------------------------	--

Evidence level	Very low
----------------	----------

Recommendation grade	Weak
----------------------	------

Perioperative oral immunonutrition (IN)

The role of IN has been investigated thoroughly over many years. Few studies specifically address IN for PD patients, and the variation in active immune-modulating nutrients administered makes interpretation difficult. A reduction in the prevalence of infectious complications is a consistent finding in patients with gastrointestinal cancer, as are beneficial effects on surrogate endpoints [levels of interleukins and C-reactive protein (CRP)] or LOSH. A reduction in mortality has not been shown. Several recently published reviews and meta-analyses [35–41] conclude that there is a benefit from perioperative and postoperative IN in patients undergoing major gastrointestinal surgery, but results remain inconsistent [42–44]. Beneficial outcomes have been shown in a systematic review of 35 trials in patients undergoing elective surgery, in which arginine-supplemented diets were associated with a significantly

reduced prevalence of infectious complications and LOSH [45]. There is also evidence to suggest that immune-modulating nutrition may be more beneficial in undernourished rather than in normally nourished patients. However, IN could be detrimental in patients with sepsis [46]. There are no trials investigating IN within ERAS care pathways.

Summary and recommendation	The balance of evidence suggests that IN for 5–7 days perioperatively should be considered because it may reduce the prevalence of infectious complications in patients undergoing major open abdominal surgery
Evidence level	Moderate
Recommendation grade	Weak

Oral bowel preparation

Mechanical bowel preparation (MBP) may lead to dehydration and offset fluid and electrolyte balance, particularly in the elderly [47]. Meta-analyses from colonic surgery have not shown clinical benefit from MBP [48, 49]. A large and recent retrospective analysis of 200 consecutive patients undergoing PD did not find any benefit of MBP to a clear liquid diet the day before surgery [50]. No trial has compared MBP to a regimen without MBP and an unrestricted diet until midnight before surgery.

Summary and recommendation	Extrapolation of data from colonic surgery and retrospective studies in PD show that MBP has no proven benefit. MBP should not be used
Evidence level	Moderate
Recommendation grade	Strong

Preoperative fasting and preoperative treatment with carbohydrates

Fasting from midnight has been standard practice in elective surgery, but is not supported by evidence [51]. Overnight fasting increases insulin resistance and discomfort after abdominal surgery [52, 53]. Guidelines recommend the intake of clear fluids up to 2 h before the induction of anaesthesia as well as a fasting period of 6 h for solids [54]. The latter recommendation has a weak scientific basis [55]. Intake of a complex clear carbohydrate-rich drink designed for preoperative use ≤ 2 h before the induction of anaesthesia has been shown to reduce hunger, thirst and anxiety, and to decrease postoperative insulin resistance [56–58]. Earlier

resumption of gut function after colorectal surgery has also been suggested [59], and an RCT including some PD patients concluded that oral carbohydrate treatment may preserve skeletal muscle mass [60]. An RCT conducted in patients undergoing cholecystectomy did not show any benefit [61]. Data on the safety and clinical benefit of preoperative carbohydrate in patients with diabetes are sparse [62, 63], and further research is warranted in this group.

Summary and recommendation	Intake of clear fluids up to 2 h before anaesthesia does not increase gastric residual volume and is recommended before elective surgery. Intake of solids should be withheld 6 h before anaesthesia. Data extrapolation from studies in major surgery suggests that preoperative oral carbohydrate treatment should be given in patients without diabetes
Evidence level	Fluid intake: High Solid intake: Low; Carbohydrate loading: Low
Recommendation grade	Fasting: Strong Carbohydrate loading: Strong

Pre-anaesthetic medication

Anxiety makes postoperative pain more difficult to control. Pre-emptive treatment of anxiety could lower pain scores and reduce the demand for opiates [64]. However, pre-induction anxiolytic medication increases postoperative sedation [65], and a meta-analysis did not demonstrate reduced postoperative pain with pre-emptive use of analgesics [66]. Short-acting anxiolytics may be helpful in some patients during placement of an epidural catheter, and experiences from day surgery suggest that cognitive function is not significantly impaired [67]. Additionally, oral fluids and a carbohydrate-rich beverage have been shown to reduce preoperative anxiety [57]. Medications for chronic pain need to be continued on the morning of surgery, and should be prescribed in the postoperative period.

Summary and recommendation	Data from studies on abdominal surgery show no evidence of clinical benefit from preoperative use of long-acting sedatives, and they should not be used routinely. Short-acting anxiolytics may be used for procedures such as insertion of epidural catheters
Evidence level	No long-acting sedatives: Moderate
Recommendation grade	Weak

Anti-thrombotic prophylaxis

Malignant disease and major surgery increase the risk of venous thromboembolism (VTE) [68]. Unfractionated and fractionated low-dose heparins are effective at preventing VTE [69]. Fractionated low-molecular-weight heparin (LMWH) is preferable in view of compliance (once-daily administration) [70]. Treatment is usually initiated 2–12 h before surgery and continued until patients are fully mobile. A meta-analysis supports continued treatment for 4 weeks after hospital discharge [71]. Concomitant use of LMWH and epidural catheters is controversial [72–75]. It has, therefore, been recommended that the catheter be inserted ≥ 12 h after a dose of LMWH, and removed ≥ 12 h after administration of LMWH [76]. The risk of an epidural or spinal haematoma is increased in patients who are also on anti-platelet drugs or oral anticoagulants [73]. Combined prophylactic modalities have been shown to be superior to pharmacological measures only in preventing VTE [77]. Mechanical intermittent pneumatic leg compression [77], and elastic stockings may be used as adjuncts in patients who are at moderate or high risk for VTE [78].

Summary and recommendation LMWH reduces the risk of thromboembolic complications. Administration should be continued for 4 weeks after hospital discharge. Concomitant use of epidural analgesia necessitates close adherence to safety guidelines. Mechanical measures should probably be added for patients at high risk

Evidence level High
Recommendation grade Strong

Antimicrobial prophylaxis and skin preparation

There is ample evidence favouring the prescription of antimicrobial prophylaxis for major abdominal procedures [79, 80]. Trials specifically targeting patients undergoing PD were not identified. Recently published studies reported or recommended prescription in a single-dose manner [80]. However, an extra dose should be provided every 3–4 h during the procedure if drugs with a short half-life are chosen [81]. Initial administration should be as near as possible to the skin incision and ≤ 1 h before the incision [79, 82]. The choice of antibiotic is dependent upon local guidelines, and should be different from the drug of choice for treatment of established infections. Skin preparation with a scrub of chlorhexidine-alcohol has recently been claimed to be superior to povidone-iodine in preventing surgical-site infections [83]. However, the difference is

likely to be very small because excellent results are obtained with povidone-iodine [84]. Alcohol-based scrubs have been reported to be used in fire-based and burn injuries [85].

Summary and recommendation Antimicrobial prophylaxis prevents surgical-site infections and should be used in a single-dose manner initiated 30–60 min before skin incision. Repeated intraoperative doses may be necessary depending on the half-life of the drug and duration of the procedure

Evidence level High
Recommendation grade Strong

Epidural analgesia

A meta-analysis showed that continuous epidural analgesia with or without opioids provided significant improvement in postoperative pain control compared with parenteral opioids in open abdominal surgery [86]. Moreover, a Cochrane review demonstrated that continuous epidural analgesia is superior to patient-controlled intravenous opioid analgesia in relieving pain ≤ 72 h after open abdominal surgery [87]. A decreased prevalence of ileus was found for epidural administration of local anaesthetic after laparotomy compared with systemic or epidural opioids in one Cochrane review [88]. With respect to complications after abdominal or thoracic surgery, a meta-analysis [89] concluded that epidural analgesia was associated with a significantly decreased risk of postoperative pneumonia, as well as an improvement in pulmonary function and arterial oxygenation. Also, the use of epidurals has been shown to reduce insulin resistance [90]. Despite the widespread use of epidural analgesia after pancreatic surgery [91], RCTs that specifically examine the outcomes of epidural analgesia after pancreatic surgery are lacking. A retrospective study comparing epidural analgesia with intravenous analgesia after PD found that patients with epidural analgesia had lower pain scores but significantly higher rates of major complications [92]. It has been suggested that thoracic epidural analgesia after PD is associated with haemodynamic instability, which might compromise enteric anastomoses, intestinal perfusion and recovery of gastrointestinal function [92]. In experimental acute pancreatitis and in sepsis, however, thoracic epidurals improved perfusion in gastrointestinal mucosal capillaries [93]. The adverse perfusion effects of epidural analgesia appear to be related to the prolonged and extended sympathetic block. This would imply that the beneficial effects of epidural analgesia can be preserved as long as the haemodynamic consequences are adequately

controlled with vasopressors [94]. Concerns about anastomoses have been raised after colorectal surgery, but one meta-analysis did not detect differences in rates of anastomotic leaks between patients receiving postoperative local anaesthetic epidurals and those receiving systemic or epidural opioids [95].

A potential drawback with epidurals is that as many as one-third of epidurals may not function satisfactorily in some centres [96, 97]. Possible reasons may be that: catheters are not located in the epidural space; the insertion level does not cover the surgical incision; the dosage of local anaesthetic and opioid are insufficient; or pump failure. For upper transverse incisions, epidural catheters should be inserted between T5 and T8 root levels. Sensory block should be tested (cold and pinprick) before induction of general anaesthesia. Efforts should be made to check the sensory block on a daily (or more frequent) basis, and the infusion should be adjusted to provide sufficient analgesia to allow mobilisation out of bed. It has been suggested that epidural analgesia should continue for ≥ 48 h and, after a successful stop-test, oral multimodal analgesia with paracetamol and non-steroidal anti-inflammatory drugs (NSAIDS)/cyclooxygenase (COX)-2 inhibitors should be commenced together with oral opioids as required. Functioning epidural catheters may be used for a longer duration if needed. Further studies are warranted to evaluate specifically the potential risks and benefits of epidural analgesia after pancreatic surgery. The use of epidurals has not been investigated for laparoscopic pancreatic resections.

Summary and recommendation	Mid-thoracic epidurals are recommended based on data from studies on major open abdominal surgery showing superior pain relief and fewer respiratory complications compared with intravenous opioids
Evidence level	Pain: High; Reduced respiratory complications: Moderate; Overall Morbidity: Low
Recommendation grade	Weak

Intravenous analgesia

Thoracic epidural anaesthesia remains the ‘gold standard’ method for major open abdominal surgery, but there are situations in which it cannot be employed. Patient-controlled analgesia (PCA) with opioids is the most common modality used as an alternative to an epidural. In a clinical trial on the implementation of a critical pathway for distal pancreatic surgery, PCA was the only analgesic modality used, but no comments were made on the impact of systemic analgesia on accelerating recovery [98].

Intravenous infusion of lidocaine has analgesic, anti-inflammatory and antihyperalgesic properties, and has been evaluated as an analgesic modality for abdominal surgery. A systematic review of 8 trials (161 patients) in which the continuous infusion of lidocaine was compared with PCA morphine for abdominal surgery, showed a decrease in the duration of ileus, LOSH, postoperative pain intensity and side effects [99]. A recent RCT in patients undergoing laparoscopic colorectal resection using the ERAS programme showed no difference in return of gastrointestinal function and LOSH between continuous infusion of lidocaine and thoracic epidural anaesthesia [100].

Summary and recommendation	Some evidence supports the use of PCA or intravenous lidocaine analgesic methods. There is insufficient information on outcome after PD
Evidence level	PCA: Very Low; I.V. Lidocaine: Moderate
Recommendation grade	Weak

Wound catheters and transversus abdominis plane (TAP) block

The efficacy of wound infusion with local anaesthetic agents as a postoperative analgesic method has been proven in a meta-analysis of different surgical procedures [101]. Conversely, a more recent meta-analysis showed that wound catheters provided no significant reduction in pain intensity (at rest or with activity) or in morphine consumption at any time after laparotomy [102]. No significant differences in the prevalence of infectious complications were found. These inconsistent results might be due to factors such as the type, concentration and dose of local anaesthetic, type of catheter, mode of delivery, or catheter location (subcutaneous or subfascial) [103]. In patients undergoing colorectal surgery, a significant opioid-sparing effect and reduction of LOSH were demonstrated when local anaesthetic was infused through a catheter positioned between the fascia and the peritoneum [104]. No significant increase in wound infections was found with the insertion of a catheter and infusion of local anaesthetics. No comparison has been made with other modalities (e.g., epidural analgesia) or in enhanced recovery programmes.

TAP blocks anaesthetise the thoracolumbar nerves (intercostal, subcostal and first lumbar), which provide sensory innervation to the anterolateral abdominal wall. The ultrasonography-guided technique for TAP blocks has been used for postoperative analgesia after abdominal surgery. A systematic analysis of 7 studies (360 patients) showed significant opioid-sparing in the postoperative period [105]. A meta-analysis of 5 RCTs (176 patients)

confirmed previous results showing improved pain relief and reduced opioid-associated side effects [106]. However, no studies have compared TAP block with other analgesic methods such as epidural analgesia or infiltration of local anaesthetic into the abdominal wound. Furthermore, no studies have used an enhanced recovery programme [107] and no studies have been conducted in patients undergoing pancreatic surgery.

The marked heterogeneity observed between studies included in the meta-analyses mentioned above would imply that further trials are needed to evaluate the potential use of wound catheters and TAP blocks in pancreatic surgery.

Summary and recommendation	Some evidence supports the use of wound catheters or TAP blocks in abdominal surgery. Results are conflicting and variable and mostly from studies in lower gastrointestinal surgery
Evidence level	Wound catheters: Moderate; TAP blocks: Moderate
Recommendation grade	Weak

Postoperative nausea and vomiting (PONV)

Data specifically addressing PONV after PD specifically have not been identified. One comparative (non-randomised) study [7] showed that an ERAS protocol with early mobilisation, metoclopramide and removal of nasogastric tube on day 1 or day 2 decreased the rate of postoperative nausea and vomiting. Until further documentation becomes available, the suggestions for patients undergoing colorectal surgery [3] should be applicable to those undergoing PD: Patients with two risk factors (female sex, non-smoking status, history of motion sickness (or PONV) and postoperative administration of opioids [108, 109]) should receive prophylaxis with dexamethasone at induction or a serotonin receptor antagonist (e.g., ondansetron, tropisetron) at the end of surgery [110]. High-risk individuals (three factors) should receive general anaesthesia with propofol and remifentanyl and no volatile anaesthetics; and dexamethasone 4–8 mg at the beginning of surgery, supplemented with serotonin receptor antagonists or droperidol [110], or 25–50 mg metoclopramide 30–60 min before the end of surgery [111]. Ondansetron can be used for prophylaxis and treatment. A possible risk of impaired anastomotic healing caused by single-dose dexamethasone or other steroids perioperatively has been addressed clinically and experimentally, but remains unclear [112–115].

Summary and recommendation	Data from the literature on gastrointestinal surgery in patients at risk of
----------------------------	---

PONV show the benefits of using different pharmacological agents depending on the patient's PONV history, type of surgery and type of anaesthesia. Multimodal intervention, during and after surgery is indicated

Evidence level	Low
Recommendation grade	Strong

Incision

There are no data comparing the types of incisions for patients undergoing PD. The authors of these recommendations are comfortable with straight transverse, curved transverse and chevron incisions, indicating that all are practical. Laparoscopic resection of the pancreatic head has been reported to be feasible [116], but its future role is uncertain.

Summary and recommendation	The choice of incision is at the surgeon's discretion, and should be of a length sufficient to ensure good exposure
----------------------------	---

Evidence level	Very Low
Recommendation grade	Strong

Avoiding hypothermia

Several meta-analyses and RCTs have demonstrated that preventing inadvertent hypothermia during major abdominal surgery (such as PD) reduces the prevalence of wound infections [117, 118], cardiac complications [118–120], bleeding and transfusion requirements [118–121], as well as the duration of post-anaesthetic recovery [122]. Furthermore, extending systemic warming in the perioperative period (2 h before and after surgery) has additional benefits [123]. Hence, the use of active cutaneous warming is highly recommended to reduce postoperative morbidity and enhance recovery. There is even evidence to suggest that circulating-water garments offer better temperature control than forced-air warming systems [124–126].

Summary and recommendation	Intraoperative hypothermia should be avoided by using cutaneous warming, i.e., forced-air or circulating-water garment systems
----------------------------	--

Evidence level	High
Recommendation grade	Strong

Postoperative glycaemic control

Postoperative hyperglycaemia in patients without diabetes is a result of acquired insulin resistance. Morbidity and mortality after major abdominal surgery have been associated with increasing levels of insulin resistance [127] and plasma glucose [128]. Such an association has also been demonstrated in pancreatic surgery [129]. Data from patients subjected to colorectal surgery within an ERAS regimen indicate that higher preoperative levels of glycated haemoglobin (HBA1c) and higher postoperative levels of glucose also predict postoperative morbidity [130].

Core elements of ERAS protocols attenuate postoperative insulin resistance and thus also lower glucose levels [131, 132]. The most obvious (of several) protocols are avoidance of preoperative fasting and oral bowel preparation; use of oral carbohydrate treatment and stimulation of early resumption of gut function by optimal fluid balance and avoidance of systemic opioids; and the reduction of the stress response by use of epidural anaesthesia.

Reducing the rate of hyperglycaemia in surgical patients in intensive-care settings has been documented to reduce the rate of complications [133–136]. Similar trials in ward settings in patients treated with modern care regimens are wanting. The target concentration for plasma glucose is controversial [137], but it seems fair to advocate that hyperglycaemia should be avoided and that this will improve outcome irrespective of the baseline level. Achieving tight glycaemic control with intravenous insulin is challenging in the ward setting because of the risk of hypoglycaemia. Glucosuria with the risk of hypovolaemia will ensue when the renal threshold is passed at >12 mmol/l [137]. This level has been used as the control regimen in seminal trials [133, 138] and should probably be regarded as a limit irrespective of settings to avoid additional disturbances in fluid balance.

Summary and recommendation	Insulin resistance and hyperglycaemia are strongly associated with postoperative morbidity and mortality. Treatment of hyperglycaemia with intravenous insulin in the intensive-care setting improves outcomes but hypoglycaemia remains a risk. Several ERAS protocol items attenuate insulin resistance and facilitate glycaemic control without the risk of hypoglycaemia. Hyperglycaemia should be avoided as far as possible without introducing the risk of hypoglycaemia
----------------------------	---

Evidence level	Low
Recommendation grade	Strong

Nasogastric intubation

There is strong evidence that routine nasogastric decompression after elective laparotomy should be avoided [139]. Fever, atelectasis and pneumonia occur more frequently in patients with a nasogastric tube than in those without [139, 140]. Bowel function returns earlier in patients if nasogastric decompression is avoided [139]. Gastro-oesophageal reflux is increased during laparotomy if nasogastric tubes are inserted [141]. The role of nasogastric tubes has not been investigated prospectively in pancreatic surgery. However, the abundant high-level evidence in other fields of abdominal surgery, including gastroduodenal surgery [139], should allow for an extrapolation to patients undergoing PD and justify a ‘no decompressive nasogastric tube’ policy. This is also supported by some series with historic controls [142, 143]. A large Norwegian RCT in patients after upper gastrointestinal and hepatopancreaticobiliary surgery (and including >80 patients who had undergone PD and were treated without routine use of a nasogastric tube) found that early oral feeding was safe and feasible [144]. This has also been corroborated by other non-randomised, fast-track implementation series in this field [5–7, 9]. In keeping with data in other areas of gastrointestinal surgery, nasogastric decompression tubes had to be replaced in ≤ 15 % of patients [6, 7, 9]. Nasogastric tubes placed during surgery (to evacuate air) should be removed before the reversal of anaesthesia. Delayed gastric emptying is a specific problem in ≈ 10 –25 % of patients after PD [6, 7, 9] and it may be necessary to insert a decompression tube in a minority of patients postoperatively.

Summary and recommendation	Pre-emptive use of nasogastric tubes postoperatively does not improve outcomes and their use is not warranted routinely
----------------------------	---

Evidence level	Moderate
Recommendation grade	Strong

Fluid balance

Patients undergoing abdominal surgery often receive excessive volumes of intravenous fluids during and in the days after surgery. This frequently exceeds actual fluid losses, resulting in a weight gain of 3–6 kg [145, 146]. Excessive overload of salt and water in the perioperative period increases postoperative complication rates and delays the return of gastrointestinal function [146–149]. This strongly suggests that near-zero fluid balance must be achieved perioperatively. Identifying the correct amount needed is a challenge that is also complicated by the use of

epidural analgesia, which causes vasodilatation and intravascular hypovolaemia with hypotension, which is often interpreted and treated as fluid depletion. The result is copious volumes of fluid administration when a vasopressor would be preferable [150]. In a meta-analysis of elective colorectal patients, intraoperative flow-guided fluid therapy with trans-oesophageal Doppler (TOD) ultrasonography to accurately assess and monitor fluid status in relation to cardiac output reduced complications and LOSH [151]. Other methods, such as lithium dilution (LiDCO) are evolving and may prove to be equivalent to TOD.

Hyperchloraemic acidosis results from infusion of 0.9 % saline. Recent studies have shown that excessive use of 0.9 % saline leads to renal oedema, reduced flow velocity in the renal artery, renal cortical tissue perfusion [152], and an overall increase in postoperative complications when compared with balanced crystalloids [153]. A recent meta-analysis [154] has suggested that postoperative complications and LOSH are significantly reduced if patients undergoing major abdominal surgery are maintained in fluid balance rather than fluid imbalance. The meta-analysis concluded that too much and too little fluid is detrimental to outcome. Although colloids produce better blood volume expansion and less interstitial space overload than crystalloids [155], there is no evidence from clinical trials and meta-analyses that colloids result in better clinical outcome than crystalloids [156]. To avoid unnecessary fluid overload, vasopressors should be considered for intra- and postoperative management of epidural-induced hypotension.

Summary and recommendation Near-zero fluid balance as well as avoiding overload of salt and water results in improved outcomes. Perioperative monitoring of stroke volume with trans-oesophageal Doppler to optimize cardiac output with fluid boluses improves outcomes. Balanced crystalloids should be preferred to 0.9 % saline

Evidence level Fluid balance: High; Oesophageal Doppler: Moderate; Balanced crystalloids vs. 0.9 % saline: Moderate

Recommendation grade Strong

Perianastomotic drains

Perianastomotic drains are believed to ameliorate the consequences of minor leaks and allow them to be treated as controlled fistulas. One RCT comparing suction drain to no drain after pancreatic cancer resection did not show significant differences in terms of mortality or overall

complication rate [157]. Moreover, patients who used these drains had a significantly greater incidence of intra-abdominal collections or fistulas (pancreatic and entero-cutaneous) [157]. A series with historic controls failed to identify any increased risk after a no-drain regimen, but this design is prone to selection bias [158]. Evaluation of early (postoperative day 3) versus late (postoperative day 5 and beyond) drain removal has been examined in an RCT [159]. Early removal of the drain in patients at low risk of pancreatic fistula (amylase value in drains <5,000 U/L at postoperative day 3) was associated with a significantly decreased rate of pancreatic fistula, abdominal and pulmonary complications. Until further data are available, a conservative approach with systematic postoperative drainage and early removal in patients at low risk of pancreatic fistula (firm pancreas, wide pancreatic duct [159–161]) is recommended. In accordance with this notion, it would seem wise to place a drain in patients with a soft pancreas and narrow duct, and leave this drain in situ slightly longer.

Summary and recommendation Early drain removal after 72 h may be advisable in patients at low risk (i.e., amylase content in drain <5,000 U/L) for developing a pancreatic fistula. There is insufficient evidence to recommend no routine use of drains routinely, but their use is based only on low-level evidence

Evidence level Early removal: High

Recommendation grade Early removal: Strong

Somatostatin analogues

Somatostatin and its synthetic analogues (e.g., octreotide) reduce splanchnic blood flow and the release of pancreatic exocrine secretion [162]. The *rationale* for its use is to reduce the risk of pancreatic anastomotic fistulas by decreasing the volume of pancreatic exocrine secretions. Several RCTs have resulted in four systemic reviews and meta-analyses that assessed the possible role of a protective effect in pancreatic surgery [163–166]. The most recent meta-analysis involved 17 trials with 1,457 patients undergoing PD and 686 undergoing distal or other resections [166]. The authors concluded that the use of somatostatin analogues reduced the crude rate of pancreatic fistulas, but that the rate of clinically significant fistulas as well as the overall major morbidity and mortality remained unchanged [166]. Subgroup analyses of the PD patients showed no significant effect of somatostatin/octreotide on any of the reported outcomes [166]. The beneficial effect of somatostatin commonly believed to be present in cases with acknowledged risk factors (soft

pancreas, small pancreatic duct) is not substantiated by the available evidence.

Summary and recommendation	Somatostatin and its analogues have no beneficial effects on outcome after PD. In general, their use is not warranted. Subgroup analyses for the variability in the texture and duct size of the pancreas are not available
Evidence level	Moderate
Recommendation grade	Strong

Urinary drainage

A meta-analysis of RCTs on urinary drainage after surgery showed that suprapubic catheterisation was superior to transurethral catheterisation [167]. Patients found suprapubic catheters more acceptable, and morbidity was reduced [167]. Most trials in the meta-analysis evaluated urinary drainage for 4–7 days. The only trial in the meta-analysis focusing specifically on hepatopancreaticobiliary surgery [168] included 82 such patients out of a total of 146. The number of patients undergoing PD was not stated. The authors found no difference in outcomes, but argued that suprapubic catheterisation is probably superior; however, the difference is likely to be small. A recent RCT with a large number of patients undergoing major surgery with thoracic epidurals found removal of transurethral catheter on postoperative day 1 to be superior in terms of infection rates and did not lead to an increased rate of re-catheterisation when compared with removal on day 3–5 [169].

Summary and recommendation	Suprapubic catheterisation is superior to transurethral catheterisation if used for >4 days. Transurethral catheters can be removed safely on postoperative day 1 or 2 unless otherwise indicated
Evidence level	High
Recommendation grade	For suprapubic: Weak; Transurethral catheter out postoperative day 1–2: Strong

Delayed gastric emptying (DGE)

DGE is a specific problem after PD occurring in \approx 10–25 % of patients [6, 7, 9, 170]. It may be necessary to insert a nasojejunal feeding tube in a minority of patients. DGE is as common after pylorus-preserving PD as after a classic Whipple's procedure [171]. In this context, DGE was less common in a fast-track group compared with a traditional care

group in one study [7]. For pylorus-preserving PDs, it has been shown that constructing the duodenojejunostomy in an antecolic (as opposed to a retro-colic) fashion results in less DGE [172]. Occasionally, DGE persists and may necessitate enteral feeding delivered beyond the gastrojejunostomy (or even parenteral nutrition). The available definition of DGE [170] is based on the assessed need for nasogastric tubes. The entity is susceptible to being over-diagnosed, and care should be taken to ensure that it does not encourage the insertion of nasogastric tubes as routine practice.

Summary and recommendation	There are no acknowledged strategies to avoid DGE. Artificial nutrition should be considered selectively in patients with DGE of long duration
Evidence level	Very low
Recommendation grade	Strong

Stimulation of bowel movement

There is no high-level evidence to support a specific motility-enhancing drug. A multimodal approach involving the use of oral laxatives such as magnesium sulphate or bisacodyl may induce early gastrointestinal transit after colonic resections [173, 174]. Some protocols for fast-track pancreatic surgery have recommended the use of laxatives postoperatively [175]. In a series of 255 pancreatic resections (almost 60 % PDs), oral administration of magnesium (200 mg/day) and lactulose in addition to metoclopramide on postoperative day 1 to support early start of normal bowel function was advocated [6]. Along with other multimodal prescriptions, the authors concluded that this protocol was associated with a low prevalence of re-admission to hospital, mortality, and morbidity rates [6]. However, no randomised trial has investigated the use of oral laxatives, so further studies are necessary. As noted above, the appropriate use of epidurals and maintaining a near-zero fluid balance are associated with an enhanced return of bowel activity after abdominal surgery [88, 146]. Chewing gum has been shown to be safe and beneficial in restoring gut activity after colorectal surgery [176–178]

Summary and recommendation	A multimodal approach with epidural and near-zero fluid balance is recommended. Oral laxatives and chewing gum given postoperatively are safe and may accelerate gastrointestinal transit
Evidence level	Laxatives: Very low; Chewing gum: Low
Recommendation grade	Weak

Postoperative artificial nutrition

Most patients tolerate normal oral intake soon after elective PD. Early oral intake in this patient group has been shown to be feasible and safe [6, 144]. A recent large multicentre RCT in patients undergoing only major upper gastrointestinal and hepatopancreaticobiliary surgery (including >80 patients undergoing PD) investigated this issue and concluded that allowing early diet is safe for these patients and that enteral tube feeding did not confer benefit [144]. This is in keeping with other reports, [179] including enteral tube feeding after other major abdominal surgery [180]. There are no data to support the idea that a surgeon-controlled stepwise increase from spoonfuls of water to a normal diet is safer than a patient-controlled routine as long as patients are informed about the potential of impaired gut function in the early postoperative period. Enteral or parenteral nutritional support will often be necessary if major complications develop. Parenteral nutrition is indicated only in those patients who cannot eat and drink normally, and who in addition cannot tolerate enteral nutrition [181]. Parenteral nutrition should be reduced as the tolerance of enteral nutritional intake increases.

Enteral tube feeding delivers artificial nutrients, but is a non-volitional intervention that bypasses the cephalic-vagal digestive reflex and carries significant risks [182, 183]. Traditionally, benefit has been shown compared with parenteral nutrition and is based on an assumption that an early- or patient-controlled oral diet is unacceptable [31]. The superiority of enteral tube feeding over an early oral diet after major abdominal surgery (including after PD), has not been documented and the opposite might well be the case (as outlined above). Oral nutritional supplementation post-hospital discharge seems appealing in a patient group known to struggle to achieve dietary goals, but evidence for a benefit is lacking [184].

Summary and recommendation	Patients should be allowed a normal diet after surgery without restrictions. They should be cautioned to begin carefully and increase intake according to tolerance over 3–4 days. Enteral tube feeding should be given only on specific indications and parenteral nutrition should not be employed routinely
Evidence level	Early diet at will: Moderate
Recommendation grade	Strong

Early and scheduled mobilisation

The relatively slow resumption of function in the stomach and gut together with significant surgical trauma leads to a

prolonged recovery period in PD patients compared with many other laparotomy patients even in the absence of major complications. Extended bed rest is associated with several unwanted effects [185, 186]. Scientific data are lacking, but the authors have observed the feasibility of written instructions for patients with detailed day-to-day targets postoperatively. This ensures autonomy and cooperation from patients. Daily progress can be monitored with diaries or with simple monitoring devices for patient activity. Analgesia must be adequate not only for rest, but also for early mobilisation.

Summary and recommendation	Patients should be mobilized actively from the morning of the first postoperative day and encouraged to meet the daily targets for mobilisation
Evidence level	Very low
Recommendation grade	Strong

Audit

Systematic audit is essential to determine clinical outcome and to establish the successful implementation and continued use of a care protocol. There are also indications that audit *per se* improves clinical results through feedback [187]. It is vital to distinguish between unsuccessful implementation and lack of desired effect from an implemented protocol if results are short of the desired quality standards. Comparison with other centres using similar protocols *via* identical tools of registration and identical definitions of key factors is needed.

Summary and recommendation	Systematic audit improves compliance and clinical outcomes
Evidence level	Low
Recommendation grade	Strong

Conclusion

ERAS[®] programmes have been strongly associated with reduced LOSH but this may not be the best indicator of the quality of functional recovery. An awareness of goals that improve safety and clinical outcomes is of greater importance. Emphasis must be placed on reducing morbidity with the introduction of standardised and appropriate enhanced recovery programmes based on best available scientific evidence.

Multimodal ERAS programmes are complex interventions that pose significant challenges to evaluation by conventional RCTs [175, 188]. The most obvious of these

challenges are standardisation of the intervention and a rapidly closing window of opportunity from ethical and practical concerns [189]. This may, to some extent, explain the relative paucity of RCTs evaluating ERAS programmes and the somewhat limited effect that has been shown on endpoints other than LOSH. In addition, interventions like these pathways are prone to show significant Hawthorne or Trial effects [190, 191]. This implies that the collateral effect on the infrastructure and management culture to implement such a comprehensive programme will have beneficial consequences in addition to those caused by the protocol items themselves or their synergistic effect. As has also been pointed out for this patient group [175], this is nevertheless a benefit related to the use of these programmes. For these reasons it may be argued that a randomised evaluation of an evidence-based ERAS protocol against traditional care may not be the way forward. Furthermore, it seems reasonable to propose that, if RCTs have proven the benefit (item by item) of two wheels, two pedals, a frame, a chain and a handle bar, then a bicycle is highly likely to be a valuable tool. Feasibility, however, must be ensured. Hence, multicentre and multinational prospective validation of a unified and comprehensive perioperative care protocol in consecutive cohorts of patients undergoing PD is warranted.

Conflict of interest The ERAS Society® receives an unrestricted grant from Nutricia Clinical Care. KL, MMEC, KS, FC, MS, RWP, KCHF, ND, MB, CHCD: Declare no conflicts of interests. JEdA-N has received travel support for lectures and meetings from Nestle and Fresenius-Kabi. DNL has received unrestricted research funding, travel grants and speaker's honoraria from Baxter Healthcare, Fresenius Kabi and BBraun. OL has served as an external Advisor to Nutricia Clinical Care, and received travel and lecture honoraria from Nutricia, Fresenius-Kabi, BBraun, Baxter and Nestle. OL previously held a patent for a preoperative carbohydrate drink formerly licensed to Nutricia.

References

1. Varadhan KK, Neal KR, Dejong CH et al (2010) The enhanced recovery after surgery (ERAS) pathway for patients undergoing major elective open colorectal surgery: a meta-analysis of randomized controlled trials. *Clin Nutr* 29:434–440
2. Fearon KC, Ljungqvist O, Von Meyenfeldt M et al (2005) Enhanced recovery after surgery: a consensus review of clinical care for patients undergoing colonic resection. *Clin Nutr* 24:466–477
3. Lassen K, Soop M, Nygren J et al (2009) Consensus review of optimal perioperative care in colorectal surgery. *Arch Surg* 144:961–969
4. Wichmann MW, Roth M, Jauch KW et al (2006) A prospective clinical feasibility study for multimodal “fast track” rehabilitation in elective pancreatic cancer surgery. *Rozhl Chir* 85:169–175
5. Kennedy EP, Rosato EL, Sauter PK et al (2007) Initiation of a critical pathway for pancreaticoduodenectomy at an academic institution—the first step in multidisciplinary team building. *J Am Coll Surg* 204:917–923
6. Berberat PO, Ingold H, Gulbinas A et al (2007) Fast track—different implications in pancreatic surgery. *J Gastrointest Surg* 11:880–887
7. Balzano G, Zerbi A, Braga M et al (2008) Fast-track recovery programme after pancreatico- duodenectomy reduces delayed gastric emptying. *Br J Surg* 95:1387–1393
8. Montiel Casado MC, Pardo SF, Rotellar SF et al (2010) Experience of a cephalic pancreatoduodenectomy fast-track program. *Cir Esp* 87:378–384
9. di Sebastiano P, Festa L, De Bonis A et al (2011) A modified fast-track program for pancreatic surgery: a prospective single-center experience. *Langenbecks Arch Surg* 396:345–351
10. Guyatt GH, Oxman AD, Vist GE et al (2008) GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 336:924–926
11. Guyatt GH, Oxman AD, Kunz R et al (2008) What is “quality of evidence” and why is it important to clinicians? *BMJ* 336:995–998
12. Guyatt GH, Oxman AD, Kunz R et al (2008) Going from evidence to recommendations. *BMJ* 336:1049–1051
13. Halaszynski TM, Juda R, Silverman DG (2004) Optimizing postoperative outcomes with efficient preoperative assessment and management. *Crit Care Med* 32:S76–S86
14. Carli F, Charlebois P, Baldini G et al (2009) An integrated multidisciplinary approach to implementation of a fast-track program for laparoscopic colorectal surgery. *Can J Anaesth* 56:837–842
15. Stergiopoulou A, Birbas K, Katostaras T et al (2007) The effect of interactive multimedia on preoperative knowledge and post-operative recovery of patients undergoing laparoscopic cholecystectomy. *Methods Inf Med* 46:406–409
16. Edward GM, Naald NV, Oort FJ et al (2011) Information gain in patients using a multimedia website with tailored information on anaesthesia. *Br J Anaesth* 106:319–324
17. Haines TP, Hill AM, Hill KD et al (2011) Patient education to prevent falls among older hospital inpatients: a randomized controlled trial. *Arch Intern Med* 171:516–524
18. Clarke HD, Timm VL, Goldberg BR et al (2011) Preoperative patient education reduces in-hospital falls after total knee arthroplasty. *Clin Orthop Relat Res* 470:244–249
19. Sewnath ME, Karsten TM, Prins MH et al (2002) A meta-analysis on the efficacy of preoperative biliary drainage for tumors causing obstructive jaundice. *Ann Surg* 236:17–27
20. Saleh MM, Norregaard P, Jorgensen HL et al (2002) Preoperative endoscopic stent placement before pancreaticoduodenectomy: a meta-analysis of the effect on morbidity and mortality. *Gastrointest Endosc* 56:529–534
21. Wang Q, Gurusamy KS, Lin H et al (2008) Preoperative biliary drainage for obstructive jaundice. *Cochrane Database Syst Rev* 3:CD005444
22. Qiu YD, Bai JL, Xu FG et al (2011) Effect of preoperative biliary drainage on malignant obstructive jaundice: a meta-analysis. *World J Gastroenterol* 17:391–396
23. Garcea G, Chee W, Ong SL et al (2010) Preoperative biliary drainage for distal obstruction: the case against revisited. *Pancreas* 39:119–126
24. van der Gaag NA, Rauws EA, van Eijck CH et al (2010) Preoperative biliary drainage for cancer of the head of the pancreas. *N Engl J Med* 362:129–137
25. Eshuis WJ, van der Gaag NA, Rauws EA et al (2010) Therapeutic delay and survival after surgery for cancer of the pancreatic head with or without preoperative biliary drainage. *Ann Surg* 252:840–849

26. Tonnesen H, Kehlet H (1999) Preoperative alcoholism and postoperative morbidity. *Br J Surg* 86:869–874
27. Tonnesen H, Rosenberg J, Nielsen HJ et al (1999) Effect of preoperative abstinence on poor postoperative outcome in alcohol misusers: randomised controlled trial. *BMJ* 318:1311–1316
28. Bluman LG, Mosca L, Newman N et al (1998) Preoperative smoking habits and postoperative pulmonary complications. *Chest* 113:883–889
29. Sorensen LT, Karlsmark T, Gottrup F (2003) Abstinence from smoking reduces incisional wound infection: a randomized controlled trial. *Ann Surg* 238:1–5
30. Lindstrom D, Sadr AO, Wladis A et al (2008) Effects of a perioperative smoking cessation intervention on postoperative complications: a randomized trial. *Ann Surg* 248:739–745
31. Goonetilleke KS, Siriwardena AK (2006) Systematic review of peri-operative nutritional supplementation in patients undergoing pancreaticoduodenectomy. *JOP* 7:5–13
32. van Stijn MF, Korkic-Halilovic I, Bakker MS et al (2012) Preoperative nutrition status and postoperative outcome in elderly general surgery patients: a systematic review. *JPEN J Parenter Enteral Nutr*. doi:10.1177/0148607112445900
33. Grotenhuis BA, Wijnhoven BP, Grune F et al (2010) Preoperative risk assessment and prevention of complications in patients with esophageal cancer. *J Surg Oncol* 101:270–278
34. Heys SD, Schofield AC, Wahle KW et al (2005) Nutrition and the surgical patient: triumphs and challenges. *Surgeon* 3: 139–144
35. Cerantola Y, Hubner M, Grass F et al (2011) Immunonutrition in gastrointestinal surgery. *Br J Surg* 98:37–48
36. Marimuthu K, Varadhan KK, Ljungqvist O et al (2012) A meta-analysis of the effect of combinations of immune modulating nutrients on outcome in patients undergoing major open gastrointestinal surgery. *Ann Surg* 255:1060–1068
37. Chen B, Zhou Y, Yang P et al (2010) Safety and efficacy of fish oil-enriched parenteral nutrition regimen on postoperative patients undergoing major abdominal surgery: a meta-analysis of randomized controlled trials. *JPEN J Parenter Enteral Nutr* 34:387–394
38. Gustafsson UO, Ljungqvist O (2011) Perioperative nutritional management in digestive tract surgery. *Curr Opin Clin Nutr Metab Care* 14:504–509
39. Marik PE, Zaloga GP (2010) Immunonutrition in high-risk surgical patients: a systematic review and analysis of the literature. *JPEN J Parenter Enteral Nutr* 34:378–386
40. Wang Y, Jiang ZM, Nolan MT et al (2010) The impact of glutamine dipeptide-supplemented parenteral nutrition on outcomes of surgical patients: a meta-analysis of randomized clinical trials. *JPEN J Parenter Enteral Nutr* 34:521–529
41. Wei C, Hua J, Bin C et al (2010) Impact of lipid emulsion containing fish oil on outcomes of surgical patients: systematic review of randomized controlled trials from Europe and Asia. *Nutrition* 26:474–481
42. Sultan J, Griffin SM, Di FF et al (2012) Randomized clinical trial of omega-3 fatty acid-supplemented enteral nutrition versus standard enteral nutrition in patients undergoing oesophagogastric cancer surgery. *Br J Surg* 99:346–355
43. Fujitani K, Tsujinaka T, Fujita J et al (2012) Prospective randomized trial of preoperative enteral immunonutrition followed by elective total gastrectomy for gastric cancer. *Br J Surg* 99:621–629
44. Mudge L, Isenring E, Jamieson GG (2011) Immunonutrition in patients undergoing esophageal cancer resection. *Dis Esophagus* 24:160–165
45. Drover JW, Dhaliwal R, Weitzel L et al (2011) Perioperative use of arginine-supplemented diets: a systematic review of the evidence. *J Am Coll Surg* 212:385–399
46. McClave SA, Martindale RG, Vanek VW et al (2009) Guidelines for the Provision and Assessment of Nutrition Support Therapy in the Adult Critically Ill Patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.). *JPEN J Parenter Enteral Nutr* 33:277–316
47. Holte K, Nielsen KG, Madsen JL et al (2004) Physiologic effects of bowel preparation. *Dis Colon Rectum* 47:1397–1402
48. Guenaga KF, Matos D, Castro AA et al (2005) Mechanical bowel preparation for elective colorectal surgery. *Cochrane Database Syst Rev* 1:CD001544
49. Cao F, Li J, Li F (2011) Mechanical bowel preparation for elective colorectal surgery: updated systematic review and meta-analysis. *Int J Colorectal Dis* 27:803–810
50. Lavu H, Kennedy EP, Mazo R et al (2010) Preoperative mechanical bowel preparation does not offer a benefit for patients who undergo pancreaticoduodenectomy. *Surgery* 148: 278–284
51. Ljungqvist O, Soreide E (2003) Preoperative fasting. *Br J Surg* 90:400–406
52. Smith I, Kranke P, Murat I et al (2011) Perioperative fasting in adults and children: guidelines from the European Society of Anaesthesiology. *Eur J Anaesthesiol* 28:556–569
53. Svanfeldt M, Thorell A, Brismar K et al (2003) Effects of 3 days of “postoperative” low caloric feeding with or without bed rest on insulin sensitivity in healthy subjects. *Clin Nutr* 22:31–38
54. American Society of Anesthesiologists Committee (2011) Practice guidelines for preoperative fasting and the use of pharmacologic agents to reduce the risk of pulmonary aspiration: application to healthy patients undergoing elective procedures: an updated report by the American Society of Anesthesiologists Committee on Standards and Practice Parameters. *Anesthesiology* 114:495–511
55. Miller M, Wishart HY, Nimmo WS (1983) Gastric contents at induction of anaesthesia. Is a 4-hour fast necessary? *Br J Anaesth* 55:1185–1188
56. Ljungqvist O, Nygren J, Thorell A (2002) Modulation of postoperative insulin resistance by pre-operative carbohydrate loading. *Proc Nutr Soc* 61:329–336
57. Hausel J, Nygren J, Lagerkranser M et al (2001) A carbohydrate-rich drink reduces preoperative discomfort in elective surgery patients. *Anesth Analg* 93:1344–1350
58. Helminen H, Viitanen H, Sajanti J (2009) Effect of preoperative intravenous carbohydrate loading on preoperative discomfort in elective surgery patients. *Eur J Anaesthesiol* 26:123–127
59. Noblett SE, Watson DS, Huong H et al (2006) Pre-operative oral carbohydrate loading in colorectal surgery: a randomized controlled trial. *Colorectal Dis* 8:563–569
60. Yuill KA, Richardson RA, Davidson HIM et al (2005) The administration of an oral carbohydrate-containing fluid prior to major elective upper-gastrointestinal surgery preserves skeletal muscle mass postoperatively—a randomised clinical trial. *Clin Nutr* 24:32–37
61. Bisgaard T, Kristiansen VB, Hjortso NC et al (2004) Randomized clinical trial comparing an oral carbohydrate beverage with placebo before laparoscopic cholecystectomy. *Br J Surg* 91:151–158
62. Gustafsson UO, Nygren J, Thorell A et al (2008) Pre-operative carbohydrate loading may be used in type 2 diabetes patients. *Acta Anaesthesiol Scand* 52:946–951
63. Breuer JP, von Dossow V, von Heymann C et al (2006) Preoperative oral carbohydrate administration to ASA III-IV patients undergoing elective cardiac surgery. *Anesth Analg* 103:1099–1108
64. Caumo W, Levandovski R, Hidalgo MP (2009) Preoperative anxiolytic effect of melatonin and clonidine on postoperative

- pain and morphine consumption in patients undergoing abdominal hysterectomy: a double-blind, randomized, placebo-controlled study. *J Pain* 10:100–108
65. Caumo W, Hidalgo MP, Schmidt AP et al (2002) Effect of preoperative anxiolysis on postoperative pain response in patients undergoing total abdominal hysterectomy. *Anaesthesia* 57: 740–746
 66. Moiniche S, Kehlet H, Dahl JB (2002) A qualitative and quantitative systematic review of preemptive analgesia for postoperative pain relief: the role of timing of analgesia. *Anesthesiology* 96:725–741
 67. Walker KJ, Smith AF (2009) Premedication for anxiety in adult day surgery. *Cochrane Database Syst Rev* 4:CD002192
 68. Spyropoulos AC, Brotman DJ, Amin AN et al (2008) Prevention of venous thromboembolism in the cancer surgery patient. *Cleve Clin J Med* 75(Suppl 3):S17–S26
 69. Clagett GP, Anderson FA Jr, Geerts W et al (1998) Prevention of venous thromboembolism. *Chest* 114:531S–560S
 70. Koch A, Bouges S, Ziegler S et al (1997) Low molecular weight heparin and unfractionated heparin in thrombosis prophylaxis after major surgical intervention: update of previous meta-analyses. *Br J Surg* 84:750–759
 71. Rasmussen MS, Jorgensen LN, Wille-Jorgensen P (2009) Prolonged thromboprophylaxis with low molecular weight heparin for abdominal or pelvic surgery. *Cochrane Database Syst Rev* 1:CD004318
 72. Horlocker TT, Wedel DJ, Benzon H et al (2003) Regional anesthesia in the anticoagulated patient: defining the risks (the second ASRA Consensus Conference on Neuraxial Anesthesia and Anticoagulation). *Reg Anesth Pain Med* 28:172–197
 73. Horlocker TT, Wedel DJ, Rowlingson JC et al (2010) Regional anesthesia in the patient receiving antithrombotic or thrombolytic therapy: American Society of Regional Anesthesia and Pain Medicine Evidence-Based Guidelines (Third Edition). *Reg Anesth Pain Med* 35:64–101
 74. Liu SS, Mulroy MF (1998) Neuraxial anesthesia and analgesia in the presence of standard heparin. *Reg Anesth Pain Med* 23:157–163
 75. Tryba M (1998) European practice guidelines: thromboembolism prophylaxis and regional anesthesia. *Reg Anesth Pain Med* 23:178–182
 76. Breivik H, Bang U, Jalonen J et al (2010) Nordic guidelines for neuraxial blocks in disturbed haemostasis from the Scandinavian Society of Anaesthesiology and Intensive Care Medicine. *Acta Anaesthesiol Scand* 54:16–41
 77. Kakkos SK, Caprini JA, Geroulakos G et al (2008) Combined intermittent pneumatic leg compression and pharmacological prophylaxis for prevention of venous thromboembolism in high-risk patients. *Cochrane Database Syst Rev* 4:CD005258
 78. Lippi G, Favaloro EJ, Cervellini G (2011) Prevention of venous thromboembolism: focus on mechanical prophylaxis. *Semin Thromb Hemost* 37:237–251
 79. Bratzler DW, Houck PM (2005) Antimicrobial prophylaxis for surgery: an advisory statement from the National Surgical Infection Prevention Project. *Am J Surg* 189:395–404
 80. Nelson RL, Glenn AM, Song F (2009) Antimicrobial prophylaxis for colorectal surgery. *Cochrane Database Syst Rev* 1:CD001181
 81. Fujita S, Saito N, Yamada T et al (2007) Randomized, multicenter trial of antibiotic prophylaxis in elective colorectal surgery: single dose vs 3 doses of a second-generation cephalosporin without metronidazole and oral antibiotics. *Arch Surg* 142:657–661
 82. Steinberg JP, Braun BI, Hellinger WC et al (2009) Timing of antimicrobial prophylaxis and the risk of surgical site infections: results from the Trial to Reduce Antimicrobial Prophylaxis Errors. *Ann Surg* 250:10–16
 83. Darouiche RO, Wall MJ Jr, Itani KM et al (2010) Chlorhexidine–alcohol versus povidone–iodine for surgical-site antisepsis. *N Engl J Med* 362:18–26
 84. Tschudin-Sutter S, Frei R, Egli-Gany D et al (2012) No risk of surgical site infections from residual bacteria after disinfection with povidone–iodine–alcohol in 1014 cases: a prospective observational study. *Ann Surg* 255:565–569
 85. Rocos B, Donaldson LJ (2012) Alcohol skin preparation causes surgical fires. *Ann R Coll Surg Engl* 94:87–89
 86. Block BM, Liu SS, Rowlingson AJ et al (2003) Efficacy of postoperative epidural analgesia: a meta-analysis. *JAMA* 290: 2455–2463
 87. Werawatganon T, Charuluxanun S (2005) Patient controlled intravenous opioid analgesia versus continuous epidural analgesia for pain after intra-abdominal surgery. *Cochrane Database Syst Rev* 1:CD004088
 88. Jorgensen H, Wetterslev J, Moiniche S et al (2000) Epidural local anaesthetics versus opioid-based analgesic regimens on postoperative gastrointestinal paralysis, PONV and pain after abdominal surgery. *Cochrane Database Syst Rev* 4:CD001893
 89. Popping DM, Elia N, Marret E et al (2008) Protective effects of epidural analgesia on pulmonary complications after abdominal and thoracic surgery: a meta-analysis. *Arch Surg* 143:990–999
 90. Uchida I, Asoh T, Shirasaka C et al (1988) Effect of epidural analgesia on postoperative insulin resistance as evaluated by insulin clamp technique. *Br J Surg* 75:557–562
 91. Bruns H, Rahbari NN, Loffler T et al (2009) Perioperative management in distal pancreatectomy: results of a survey in 23 European participating centres of the DISPACT trial and a review of literature. *Trials* 10:58
 92. Pratt WB, Steinbrook RA, Maithel SK et al (2008) Epidural analgesia for pancreatoduodenectomy: a critical appraisal. *J Gastrointest Surg* 12:1207–1220
 93. Daudel F, Freise H, Westphal M et al (2007) Continuous thoracic epidural anesthesia improves gut mucosal microcirculation in rats with sepsis. *Shock* 28:610–614
 94. Hildebrand LB, Koepfli E, Kimberger O et al (2011) Hypotension during fluid-restricted abdominal surgery: effects of norepinephrine treatment on regional and microcirculatory blood flow in the intestinal tract. *Anesthesiology* 114:557–564
 95. Holte K, Kehlet H (2001) Epidural analgesia and risk of anastomotic leakage. *Reg Anesth Pain Med* 26:111–117
 96. McLeod G, Davies H, Munnoch N et al (2001) Postoperative pain relief using thoracic epidural analgesia: outstanding success and disappointing failures. *Anaesthesia* 56:75–81
 97. Burstal R, Wegener F, Hayes C et al (1998) Epidural analgesia: prospective audit of 1062 patients. *Anaesth Intensive Care* 26:165–172
 98. Kennedy EP, Grenda TR, Sauter PK et al (2009) Implementation of a critical pathway for distal pancreatectomy at an academic institution. *J Gastrointest Surg* 13:938–944
 99. Marret E, Rolin M, Beaussier M et al (2008) Meta-analysis of intravenous lidocaine and postoperative recovery after abdominal surgery. *Br J Surg* 95:1331–1338
 100. Wongyingsinn M, Baldini G, Charlebois P et al (2011) Intravenous lidocaine versus thoracic epidural analgesia: a randomized controlled trial in patients undergoing laparoscopic colorectal surgery using an enhanced recovery program. *Reg Anesth Pain Med* 36:241–248
 101. Liu SS, Richman JM, Thirlby RC et al (2006) Efficacy of continuous wound catheters delivering local anesthetic for postoperative analgesia: a quantitative and qualitative systematic review of randomized controlled trials. *J Am Coll Surg* 203:914–932

102. Gupta A, Favaio S, Perniola A et al (2011) A meta-analysis of the efficacy of wound catheters for post-operative pain management. *Acta Anaesthesiol Scand* 55:785–796
103. Yndgaard S, Holst P, Bjerre-Jepsen K et al (1994) Subcutaneously versus subfascially administered lidocaine in pain treatment after inguinal herniotomy. *Anesth Analg* 79:324–327
104. Beaussier M, El'Ayoubi H, Schiffer E et al (2007) Continuous preperitoneal infusion of ropivacaine provides effective analgesia and accelerates recovery after colorectal surgery: a randomized, double-blind, placebo-controlled study. *Anesthesiology* 107:461–468
105. Petersen PL, Mathiesen O, Torup H et al (2010) The transversus abdominis plane block: a valuable option for postoperative analgesia? A topical review. *Acta Anaesthesiol Scand* 54:529–535
106. Siddiqui MR, Sajid MS, Uncles DR et al (2011) A meta-analysis on the clinical effectiveness of transversus abdominis plane block. *J Clin Anesth* 23:7–14
107. Charlton S, Cyna AM, Middleton P et al (2010) Perioperative transversus abdominis plane (TAP) blocks for analgesia after abdominal surgery. *Cochrane Database Syst Rev* 12:CD007705
108. Apfel CC, Kranke P, Eberhart LH et al (2002) Comparison of predictive models for postoperative nausea and vomiting. *Br J Anaesth* 88:234–240
109. Rusch D, Eberhart L, Biedler A et al (2005) Prospective application of a simplified risk score to prevent postoperative nausea and vomiting. *Can J Anaesth* 52:478–484
110. Carlisle JB, Stevenson CA (2006) Drugs for preventing postoperative nausea and vomiting. *Cochrane Database Syst Rev* 3:CD004125
111. Wallenborn J, Gelbrich G, Bulst D et al (2006) Prevention of postoperative nausea and vomiting by metoclopramide combined with dexamethasone: randomised double blind multicentre trial. *BMJ* 333:324
112. Polat A, Nayci A, Polat G et al (2002) Dexamethasone down-regulates endothelial expression of intercellular adhesion molecule and impairs the healing of bowel anastomoses. *Eur J Surg* 168:500–506
113. Engelman E, Maeyens C (2010) Effect of preoperative single-dose corticosteroid administration on postoperative morbidity following esophagectomy. *J Gastrointest Surg* 14:788–804
114. De OG Jr, Almeida MD, Benzon HT et al (2011) Perioperative single dose systemic dexamethasone for postoperative pain: a meta-analysis of randomized controlled trials. *Anesthesiology* 115:575–588
115. Eubanks TR, Greenberg JJ, Dobrin PB et al (1997) The effects of different corticosteroids on the healing colon anastomosis and cecum in a rat model. *Am Surg* 63:266–269
116. Zureikat AH, Breaux JA, Steel JL et al (2011) Can laparoscopic pancreaticoduodenectomy be safely implemented? *J Gastrointest Surg* 15:1151–1157
117. Kurz A, Sessler DI, Lenhardt R (1996) Perioperative normothermia to reduce the incidence of surgical-wound infection and shorten hospitalization. Study of Wound Infection and Temperature Group. *N Engl J Med* 334:1209–1215
118. Scott EM, Buckland R (2006) A systematic review of intraoperative warming to prevent postoperative complications. *AORN J* 83:1090–2013
119. Frank SM, Fleisher LA, Breslow MJ et al (1997) Perioperative maintenance of normothermia reduces the incidence of morbid cardiac events. A randomized clinical trial. *JAMA* 277:1127–1134
120. Neshar N, Zisman E, Wolf T et al (2003) Strict thermoregulation attenuates myocardial injury during coronary artery bypass graft surgery as reflected by reduced levels of cardiac-specific troponin I. *Anesth Analg* 96:328–335
121. Rajagopalan S, Mascha E, Na J et al (2008) The effects of mild perioperative hypothermia on blood loss and transfusion requirement. *Anesthesiology* 108:71–77
122. Lenhardt R, Marker E, Goll V et al (1997) Mild intraoperative hypothermia prolongs postanesthetic recovery. *Anesthesiology* 87:1318–1323
123. Wong PF, Kumar S, Bohra A et al (2007) Randomized clinical trial of perioperative systemic warming in major elective abdominal surgery. *Br J Surg* 94:421–426
124. Galvao CM, Liang Y, Clark AM (2010) Effectiveness of cutaneous warming systems on temperature control: meta-analysis. *J Adv Nurs* 66:1196–1206
125. Taguchi A, Ratnaraj J, Kabon B et al (2004) Effects of a circulating-water garment and forced-air warming on body heat content and core temperature. *Anesthesiology* 100:1058–1064
126. Perez-Protto S, Sessler DI, Reynolds LF et al (2010) Circulating-water garment or the combination of a circulating-water mattress and forced-air cover to maintain core temperature during major upper-abdominal surgery. *Br J Anaesth* 105:466–470
127. Sato H, Carvalho G, Sato T et al (2010) The association of preoperative glycemic control, intraoperative insulin sensitivity, and outcomes after cardiac surgery. *J Clin Endocrinol Metab* 95:4338–4344
128. Jackson RS, Amdur RL, White JC et al (2012) Hyperglycemia is associated with increased risk of morbidity and mortality after colectomy for cancer. *J Am Coll Surg* 214:68–80
129. Eshuis WJ, Hermanides J, van Dalen JW et al (2011) Early postoperative hyperglycemia is associated with postoperative complications after pancreaticoduodenectomy. *Ann Surg* 253:739–744
130. Gustafsson UO, Thorell A, Soop M et al (2009) Haemoglobin A1c as a predictor of postoperative hyperglycaemia and complications after major colorectal surgery. *Br J Surg* 96:1358–1364
131. Ljungqvist O (2010) Insulin resistance and outcomes in surgery. *J Clin Endocrinol Metab* 95:4217–4219
132. Ljungqvist O (2012) Jonathan E. Rhoads lecture 2011: insulin resistance and enhanced recovery after surgery. *JPEN J Parenter Enteral Nutr* 36:389–398
133. van den Berghe G, Wouters P, Weekers F et al (2001) Intensive insulin therapy in the critically ill patients. *N Engl J Med* 345:1359–1367
134. Finfer S, Chittock DR, Su SY et al (2009) Intensive versus conventional glucose control in critically ill patients. *N Engl J Med* 360:1283–1297
135. Furnary AP, Zerr KJ, Grunkemeier GL et al (1999) Continuous intravenous insulin infusion reduces the incidence of deep sternal wound infection in diabetic patients after cardiac surgical procedures. *Ann Thorac Surg* 67:352–360
136. Krinsley JS (2004) Effect of an intensive glucose management protocol on the mortality of critically ill adult patients. *Mayo Clin Proc* 79:992–1000
137. van den BG, Schetz M, Vlasselaers D et al (2009) Clinical review: intensive insulin therapy in critically ill patients: NICE-SUGAR or Leuven blood glucose target? *J Clin Endocrinol Metab* 94:3163–3170
138. van den BG, Wilmer A, Hermans G et al (2006) Intensive insulin therapy in the medical ICU. *N Engl J Med* 354:449–461
139. Nelson R, Edwards S, Tse B (2007) Prophylactic nasogastric decompression after abdominal surgery. *Cochrane Database Syst Rev* 3:CD004929
140. Cheatham ML, Chapman WC, Key SP et al (1995) A meta-analysis of selective versus routine nasogastric decompression after elective laparotomy. *Ann Surg* 221:469–476

141. Manning BJ, Winter DC, McGreal G et al (2001) Nasogastric intubation causes gastroesophageal reflux in patients undergoing elective laparotomy. *Surgery* 130:788–791
142. Fisher WE, Hodges SE, Cruz G et al (2011) Routine nasogastric suction may be unnecessary after a pancreatic resection. *HPB (Oxford)* 13:792–796
143. Roland CL, Mansour JC, Schwarz RE (2012) Routine nasogastric decompression is unnecessary after pancreatic resections. *Arch Surg* 147:287–289
144. Lassen K, Kjæve J, Fetveit T et al (2008) Allowing normal food at will after major upper gastrointestinal surgery does not increase morbidity: a randomized multicenter trial. *Ann Surg* 247:721–729
145. Tambyraja AL, Sengupta F, MacGregor AB et al (2004) Patterns and clinical outcomes associated with routine intravenous sodium and fluid administration after colorectal resection. *World J Surg* 28:1046–1051. doi:10.1007/s00268-004-7383-7
146. Lobo DN, Bostock KA, Neal KR et al (2002) Effect of salt and water balance on recovery of gastrointestinal function after elective colonic resection: a randomised controlled trial. *Lancet* 359:1812–1818
147. Lobo DN (2009) Fluid overload and surgical outcome: another piece in the jigsaw. *Ann Surg* 249:186–188
148. Chowdhury AH, Lobo DN (2011) Fluids and gastrointestinal function. *Curr Opin Clin Nutr Metab Care* 14:469–476
149. Brandstrup B, Tonnesen H, Beier-Holgersen R et al (2003) Effects of intravenous fluid restriction on postoperative complications: comparison of two perioperative fluid regimens: a randomized assessor-blinded multicenter trial. *Ann Surg* 238:641–648
150. Holte K, Foss NB, Svensen C et al (2004) Epidural anesthesia, hypotension, and changes in intravascular volume. *Anesthesiology* 100:281–286
151. Abbas SM, Hill AG (2008) Systematic review of the literature for the use of oesophageal Doppler monitor for fluid replacement in major abdominal surgery. *Anaesthesia* 63:44–51
152. Chowdhury A, Cox E, Francis S et al (2012) A randomized, controlled, double-blind crossover study on the effects of 2-liters infusions of 0.9 % saline and Plasma-Lyte 148 on renal blood flow velocity and renal cortical tissue perfusion in healthy volunteers. *Ann Surg* 256:18–24
153. Shaw AD, Bagshaw SM, Goldstein SL et al (2012) Major complications, mortality and resource utilization after open abdominal surgery: 0.9 % saline compared to Plasma-lyte. *Ann Surg* 255:821–829
154. Varadhan KK, Lobo DN (2010) A meta-analysis of randomised controlled trials of intravenous fluid therapy in major elective open abdominal surgery: getting the balance right. *Proc Nutr Soc* 69:488–498
155. Lobo DN, Stanga Z, Aloysius MM et al (2010) Effect of volume loading with 1 liter intravenous infusions of 0.9 % saline, 4 % succinylated gelatine (Gelofusine) and 6 % hydroxyethyl starch (Voluven) on blood volume and endocrine responses: a randomized, three-way crossover study in healthy volunteers. *Crit Care Med* 38:464–470
156. Senagore AJ, Emery T, Luchtefeld M et al (2009) Fluid management for laparoscopic colectomy: a prospective, randomized assessment of goal-directed administration of balanced salt solution or hetastarch coupled with an enhanced recovery program. *Dis Colon Rectum* 52:1935–1940
157. Conlon KC, Labow D, Leung D et al (2001) Prospective randomized clinical trial of the value of intraperitoneal drainage after pancreatic resection. *Ann Surg* 234:487–493
158. Fisher WE, Hodges SE, Silberfein EJ et al (2011) Pancreatic resection without routine intraperitoneal drainage. *HPB (Oxford)* 13:503–510
159. Bassi C, Molinari E, Malleo G et al (2010) Early versus late drain removal after standard pancreatic resections: results of a prospective randomized trial. *Ann Surg* 252:207–214
160. Pratt WB, Callery MP, Vollmer CM Jr (2008) Risk prediction for development of pancreatic fistula using the ISGPF classification scheme. *World J Surg* 32:419–428. doi:10.1007/s00268-007-9388-5
161. Kawai M, Kondo S, Yamaue H et al (2011) Predictive risk factors for clinically relevant pancreatic fistula analyzed in 1,239 patients with pancreaticoduodenectomy: multicenter data collection as a project study of pancreatic surgery by the Japanese Society of Hepato-Biliary-Pancreatic Surgery. *J Hepatobiliary Pancreat Sci* 18:601–608
162. Suc B, Msika S, Piccinini M et al (2004) Octreotide in the prevention of intra-abdominal complications following elective pancreatic resection: a prospective, multicenter randomized controlled trial. *Arch Surg* 139:288–294
163. Li-Ling J, Irving M (2001) Somatostatin and octreotide in the prevention of postoperative pancreatic complications and the treatment of enterocutaneous pancreatic fistulas: a systematic review of randomized controlled trials. *Br J Surg* 88:190–199
164. Connor S, Alexakis N, Garden OJ et al (2005) Meta-analysis of the value of somatostatin and its analogues in reducing complications associated with pancreatic surgery. *Br J Surg* 92:1059–1067
165. Alghamdi AA, Jawas AM, Hart RS (2007) Use of octreotide for the prevention of pancreatic fistula after elective pancreatic surgery: a systematic review and meta-analysis. *Can J Surg* 50:459–466
166. Koti RS, Gurusamy KS, Fusai G et al (2010) Meta-analysis of randomized controlled trials on the effectiveness of somatostatin analogues for pancreatic surgery: a Cochrane review. *HPB (Oxford)* 12:155–165
167. McPhail MJ, Abu-Hilal M, Johnson CD (2006) A meta-analysis comparing suprapubic and transurethral catheterization for bladder drainage after abdominal surgery. *Br J Surg* 93:1038–1044
168. Baan AH, Vermeulen H, van der MJ et al (2003) The effect of suprapubic catheterization versus transurethral catheterization after abdominal surgery on urinary tract infection: a randomized controlled trial. *Dig Surg* 20:290–295
169. Zaouter C, Kaneva P, Carli F (2009) Less urinary tract infection by earlier removal of bladder catheter in surgical patients receiving thoracic epidural analgesia. *Reg Anesth Pain Med* 34:542–548
170. Wente MN, Bassi C, Dervenis C et al (2007) Delayed gastric emptying (DGE) after pancreatic surgery: a suggested definition by the International Study Group of Pancreatic Surgery (ISGPS). *Surgery* 142:761–768
171. Diener MK, Fitzmaurice C, Schwarzer G et al (2011) Pylorus-preserving pancreaticoduodenectomy (pp Whipple) versus pancreaticoduodenectomy (classic Whipple) for surgical treatment of periampullary and pancreatic carcinoma. *Cochrane Database Syst Rev* 5:CD006053
172. Tani M, Terasawa H, Kawai M et al (2006) Improvement of delayed gastric emptying in pylorus-preserving pancreaticoduodenectomy: results of a prospective, randomized, controlled trial. *Ann Surg* 243:316–320
173. Basse L, Madsen JL, Kehlet H (2001) Normal gastrointestinal transit after colonic resection using epidural analgesia, enforced oral nutrition and laxative. *Br J Surg* 88:1498–1500
174. Zingg U, Miskovic D, Pasternak I et al (2008) Effect of bisacodyl on postoperative bowel motility in elective colorectal surgery: a prospective, randomized trial. *Int J Colorectal Dis* 23:1175–1183

175. Ypsilantis E, Praseedom RK (2009) Current status of fast-track recovery pathways in pancreatic surgery. *JOP* 10:646–650
176. Noble EJ, Harris R, Hosie KB et al (2009) Gum chewing reduces postoperative ileus? A systematic review and meta-analysis. *Int J Surg* 7:100–105
177. Vasquez W, Hernandez AV, Garcia-Sabrido JL (2009) Is gum chewing useful for ileus after elective colorectal surgery? A systematic review and meta-analysis of randomized clinical trials. *J Gastrointest Surg* 13:649–656
178. de Castro SM, van den Esschert JW, van Heek NT et al (2008) A systematic review of the efficacy of gum chewing for the amelioration of postoperative ileus. *Dig Surg* 25:39–45
179. Yermilov I, Jain S, Sekeris E et al (2009) Utilization of parenteral nutrition following pancreaticoduodenectomy: is routine jejunostomy tube placement warranted? *Dig Dis Sci* 54:1582–1588
180. Koretz RL (2009) Enteral nutrition: a hard look at some soft evidence. *Nutr Clin Pract* 24:316–324
181. Gianotti L, Meier R, Lobo DN et al (2009) ESPEN Guidelines on Parenteral Nutrition: pancreas. *Clin Nutr* 28:428–435
182. Han-Geurts IJ, Verhoef C, Tilanus HW (2004) Relaparotomy following complications of feeding jejunostomy in esophageal surgery. *Dig Surg* 21:192–196
183. Lobo DN, Williams RN, Welch NT et al (2006) Early postoperative jejunostomy feeding with an immune modulating diet in patients undergoing resectional surgery for upper gastrointestinal cancer: a prospective, randomized, controlled, double-blind study. *Clin Nutr* 25:716–726
184. Lidder PG, Lewis S, Duxbury M et al (2009) Systematic review of postdischarge oral nutritional supplementation in patients undergoing GI surgery. *Nutr Clin Pract* 24:388–394
185. Kehlet H, Wilmore DW (2002) Multimodal strategies to improve surgical outcome. *Am J Surg* 183:630–641
186. Convertino VA (1997) Cardiovascular consequences of bed rest: effect on maximal oxygen uptake. *Med Sci Sports Exerc* 29:191–196
187. Jamtvedt G, Young JM, Kristoffersen DT et al (2006) Audit and feedback: effects on professional practice and health care outcomes. *Cochrane Database Syst Rev* 19:CD000259
188. Campbell M, Fitzpatrick R, Haines A et al (2000) Framework for design and evaluation of complex interventions to improve health. *BMJ* 321:694–696
189. Lassen K, Høye A, Myrmet T (2012) Randomised trials in surgery: the burden of evidence. *Rev Recent Clin Trials* 7:244–248
190. Franke RH, Kaul JD (1978) The Hawthorne experiments: first statistical interpretation. *Am Soc Rev* 43:623–643
191. McCarney R, Warner J, Iliffe S et al (2007) The Hawthorne Effect: a randomised, controlled trial. *BMC Med Res Methodol* 7:30